## LYCOPERDACEAE—GASTEROMYCETES

## AUTHOR CITATION AND PUBLICATION DATE OF THE NAME CALVATIA CRANIIFORMIS

In the course of our taxonomic studies on the genus *Calvatia* Fr. emend. Morgan in southern Africa, we occasionally come across interesting nomenclatural inconsistencies. An important and noteworthy case addressed in this contribution concerns the origin of the name *Calvatia craniformis*, the type of the genus *Calvatia* Fr., a conserved generic name. Originally described as *Bovista craniformis* by Von Schweinitz (1832), this taxon has subsequently been treated as a member of *Calvatia*.

Throughout the literature, Fries (1849: 442), where the genus *Calvatia* was first established, is consistently cited also as the place of valid publication of the combination *Calvatia craniiformis*. In terms of ICBN Article 33.1 (Greuter *et al.* 2000: 56), however, that is incorrect, since, when Fries (1849) established the genus *Calvatia*, he merely remarked: '*Hujus loci* Bovista cranif. *Schw.*', and nowhere in the original publication did he actually definitely associate the specific epithet *craniiformis* with the genus name *Calvatia* or its abbreviation as required by Art. 33.1. Two similar cases are discussed in Art. 33.1, Ex. 2, which make it very clear that the name *Calvatia craniiformis* was not validly published in Fries (1849).

As far as we could ascertain, the combination *Calvatia craniiformis* was first validly published by De Toni (1888: 106) in Saccardo's *Sylloge fungorum* 7. Although De Toni also ascribed this name to Fries (1849), it should be cited either as *Calvatia craniiformis* (Schwein.) Fr. ex De Toni or merely as *Calvatia craniiformis* (Schwein.) Fr. as has been the exclusive practice in the past.

## ACKNOWLEDGEMENTS

The hospitality extended to the senior author by Dr D.N. Pegler and staff of the Mycological Herbarium at Kew during a brief study visit to that institution is acknowledged with appreciation. Funding granted by the Peninsula Technikon research committee is also acknowledged with gratitude.

## REFERENCES

- DE TONI, J.B. 1888. Nidulariaceae, Lycoperdaceae et Hymenogastraceae. In P.A. Saccardo, Sylloge fungorum omnium hucusque cognitorum 7,1: 28–180. Sumptibus auctoris typis seminarii, Patavii.
- FRIES, E.M. 1849. Summa vegetabilium Scandinaviae. Sectio Posterior. Typographia Academica, Uppsala.
- GREUTER, W., MCNEILL, J., BARRIE, F.R., BURDET, H.M., DEMOULIN, V., FILGUEIRAS, T.S., NICOLSON, D.H., SILVA, P.C., SKOG, J.E., TREHANE, P., TURLAND, N.J. & HAWKSWORTH, D.L. 2000. International Code of Botanical Nomenclature (Saint Louis Code). *Regnum vegetabile* 138. Koeltz Scientific Books, Königstein.
- VON SCHWEINITZ, L.D. 1832. Synopsis fungorum in America boreali media degentium. *Transactions of the American Philo*sophical Society held at Philadelphia for promoting useful knowledge n.s. 4: 141–316 (plus plate XIX & legend).

J.C. COETZEE\* and A.E. VAN WYK\*\*

<sup>\*</sup> Department of Horticulture and Food Technology, Peninsula Technikon, P.O. Box 1906, 7535 Bellville.

<sup>\*\*</sup> H.G.W.J. Schweickerdt Herbarium, Department of Botany, University of Pretoria, 0001 Pretoria.